



Associations between intestinal lesions and detection of *Clostridium perfringens* type A or beta-2 toxin in neonatal piglets with diarrhoea

Kongsted, Hanne; Jonach, Beata Renata; Kokotovic, Branko; Nielsen, J. P.

Publication date:
2013

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Kongsted, H., Jonach, B. R., Kokotovic, B., & Nielsen, J. P. (2013). *Associations between intestinal lesions and detection of *Clostridium perfringens* type A or beta-2 toxin in neonatal piglets with diarrhoea*. Poster session presented at International Meeting on Emerging Diseases and Surveillance (IMED 2013), Vienna, Austria.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Associations between intestinal lesions and detection of *Clostridium perfringens* type A or beta-2 toxin in neonatal piglets with diarrhoea

Kongsted, H.¹, Jonach, B.², Kokotovic, B.², Nielsen, J.P.³
¹Pig Research Centre, [Danish Agriculture & Food Council](#), Denmark. hko@lf.dk
²National Veterinary Institute, Technical University of Denmark, Denmark
³Department of large animal sciences, University of Copenhagen, Denmark

Objective: To evaluate associations between gross and histopathological lesions and the presence of *Clostridium perfringens* type A (*CpA*) and beta-2 toxin in piglets from 4 herds with outbreaks of diarrhoea.


Methods: Pathological examinations on 51 diarrhoeic piglets aged 3-7 days were carried out. *CpA* was cultured and typed by PCR. A *Clostridium perfringens* spp targeted probe was used for flourescent in situ hybridization (FISH) and detection of beta-2 toxin in intestinal contents was performed by enzyme-linked immunosorbent assay (ELISA).

Results: Piglets without intestinal lesions had a significantly ($P < 0.05$) higher prevalence of *CpA* and beta-2 toxin (Table 1).

Fig.1: Flaccidity of small intestine. The intestine is thin-walled and flaccid throughout its length.



Fig. 2: FISH positive ileal mucosa. Double hybridization for Domain bacteria (green) and *Cl. perfringens* (red). Moderate amounts of *Cl. perfringens* cells are seen.



	<i>CpA</i> positive samples				Beta-2 positive samples	
Intestinal lesion	Culture (n=51)		FISH# (n=51)		ELISA (n=45)	
	Lp*	Lnp*	Lp	Lnp	Lp	Lnp
Flaccidity of small intestine	24%	64%				
Flaccidity of large intestine	19%	54%			32%	83%
Villous atrophy	22%	58%				
Small intestinal epithelial lesions			60%	90%	25%	65%

Table 1. Associations between intestinal lesions and detection of *CpA* and beta-2 toxin. Only statistically significant asociations (Fisher’s exact test, $\alpha=0.05$) are shown. # Flourescent in situ hybridization using a *Clostridium perfringens* spp targeted probe. * Lp: Lesion present. Lnp: Lesion not present.

Conclusion: Demonstration of *CpA* or Beta-2 toxin was associated with absence of intestinal lesions.